



Unit

laxseed
cultural
March,

tatum).
dustry,
ral Re

armers'
nomics.

in the
p. 887.
cultural
eration.

gricul-
eau of

es -and
eau of
eering,
1944.

nd In-
ports,
as A.

al Ex-
ruary,

ustrial
Voca-
chigan
5.

Loss.
tation,
rolina,
ervice,

ltural
rginia.

. 159.
rticul-
Mich.

Ended
iment

Cent
n. 543.
orgia.

alley.
xperi-
ation,

Office,

001



68 558a

JUNE 1945

SOIL CONSERVATION

OFFICIAL ORGAN OF THE SOIL CONSERVATION SERVICE

UNITED STATES DEPARTMENT OF AGRICULTURE, WASHINGTON, D. C.

SOIL CONSERVATION

CLAUDE R. WICKARD
SECRETARY OF AGRICULTURE

MARVIN JONES
WAR FOOD ADMINISTRATOR

HUGH H. BENNETT
CHIEF, SOIL CONSERVATION SERVICE

ISSUED MONTHLY BY SOIL CONSERVATION SERVICE, U. S. DEPT. OF AGRICULTURE, WASHINGTON, D. C.

VOL. X—No. 12

JUNE • 1945

WELLINGTON BRINK EDITOR

CONTENTS

	Page
SHELTERBELTS IN THE GREAT PLAINS. By John W. Keller.....	259
OLD WEST LIVES AGAIN IN LU PROJECTS. By O. E. McConnell	264
FIGHTING FERN WITHOUT FIRE. By Albert Arnst.....	268
GROWING PEANUTS THE CONSERVATION WAY. By W. M. Nixon	271
IMPROVE GRASSLAND FOR BETTER LIVESTOCK. By R. H. Lush	272
WATER STORAGE IN THE SURFACE SOIL. By O. R. Neal	274
ON PLANTING IN UNPLOWED STUBBLE. A translation by D. B. Krimgold	277
THE FARMER AND THE REST OF US. Reviewed by W. A. Albrecht	279
REFERENCE LIST. Compiled by William L. Robey.....	280

Front Cover: Betty Mueller with an armful of oats, on the John Mueller farm near Winona, Minn., July 1944. The grain is grown there in contour strips, of course. Photographer: W. H. Lathrop.

SOIL CONSERVATION is issued monthly by SOIL CONSERVATION SERVICE of the United States Department of Agriculture, Washington, D. C. The matter contained herein is published by direction of the Secretary of Agriculture as administrative information required for proper transaction of the public business, with the approval of the Director of the Budget. SOIL CONSERVATION seeks to supply to workers of the Department of Agriculture engaged in soil conservation activities, information of special help to them in the performance of their duties. Copies may be obtained from the Superintendent of Documents, Government Printing Office, Washington 25, D. C., 10 cents a copy, or by subscription at the rate of \$1.00 per year, domestic; \$1.50 per year, foreign. Postage stamps, will not be accepted in payment.

Part of
Even in
farmst

Shelterbelts in the Great Plains



Shelterbelts clear to the skyline. An area south of Mitchell, S. D.

By JOHN W. KELLER

In the Great Plains prolonged droughts and strong winds have intermittently harassed the rural population and made agriculture insecure. These hazards can not be eliminated entirely and any factors that aim to ameliorate them and temper their devastating effects are readily grasped by farmers and ranchers.

The planting of trees long has been recognized as a practical means of checking the effects of strong winds and the many harmful results to agriculture that accompany them. The purpose of belts of trees in the Great Plains is to deflect and slow up winds near the ground. Where this can be accomplished, it will temper the hot, dry winds of summer and the cold winds of winter, check the devastating dust storms which carry away the top soil, prevent excessive evaporation, and retain soil moisture for the production of agricultural crops.

Shelterbelts are now one of many conservation practices useful in the program of soil conservation districts, particularly in the Great Plains region.

History

Tree planting to conserve moisture for crops has been successful in Europe and, coming closer to home, in Canada as well.

EDITOR'S NOTE.—The author is assistant chief, Forestry Division, Soil Conservation Service, Washington, D. C.



Part of a 30-acre planting in the Park River area, N. D. Even in 1939 when this picture was made windbreaks and farmstead planting had vastly increased the value of the land.

In Eastern Russia shelterbelts have been planted in the steppes from the Black Sea to North of the Caspian on land that in many respects bears close resemblance to our own prairie region. The results of established shelterbelts in Russia were so favorable that the planting of over 800,000 acres of shelterbelts was included in the Soviet second 5-year plan, of which we heard so much before World War II.

In Hungary the climate, soils and precipitation southeast of Budapest are quite similar to our prairie region. Here the planting of shelterbelts for the protection of orchards, vineyards and cultivated fields has been common for many years. Legislation in Hungary in 1923 provided for a definite system of shelterbelts which were to be planted by the farmer who in return would receive planting stock at nominal prices and a reduction of taxation. These shelterbelts are reported to reduce evaporation of soil moisture, promote the formation of dew, retain litter and snow, stabilize sandy soils, protect flowering, hasten the ripening of fruit, decrease breakage of fruit trees, decrease premature fruit dropping and reduce the dust. These beneficial effects are attributed largely to a reduction of wind velocity.

In Denmark shelterbelts and hedges of trees and shrubs have proved to be the most satisfactory means of reducing high wind velocity and attendant damage. Strong winds from the North Sea in Jutland gradually buried under drifting sand a large area of once fertile agricultural soil. It is claimed that these strong winds destroy gardens, damage plants and buildings, dry out sod and whip the grain in the fields if unprotected by shelterbelts. The Danes attribute to shelterbelts favorable local effects upon air and soil temperatures, wind velocity, evaporation, relative humidity of air in local areas, and increase yields up to 30 percent for some crops.

In the prairie region in Southwestern Manitoba and portions of Saskatchewan and Alberta the soil topography and climate are quite similar to those in our Great Plains. Governmental aid and encouragement to shelterbelt plantings have assisted the development of planting methods and the testing of tree species. Early settlers in Canada planted thousands of trees about their homesteads but failures were general because of unsuitable species and lack of experience in tree culture under those rigorous conditions. In 1901 the Dominion government established experiment stations and a system of cooperative planting was begun whereby farmers were supplied with plant-

ing stock free of charge if they would carefully follow instructions in planting, cultivating and caring for the trees. Under this system the plantings are reported to be generally successful.

The beneficial effect of trees was recognized in the United States at an early date and legislation has been enacted through the years in an attempt to compensate for land misuse. In the United States the Great Plains comprise about 30 percent of the continental United States. As civilization traveled westward and increasingly large areas of sod were broken, frequent droughts and dust storms resulted in increasing economic loss and human discomfort and suffering. Tree planting was one of the first measures to relieve this distress. The Timber Culture Act of 1873, the Kincaid distribution in the agricultural appropriation act of 1911, the establishment of experiment stations in 1913, the Clarke-McNary Act of 1924 and numerous State laws encouraged tree planting. Unfortunately many of the early plantings not only received little or no care, but they were severely handicapped by heavy grazing. However, a few of them stand today as monuments to the foresight and painstaking care of the early settlers.

The planting of shelterbelts in the Great Plains is advisable wherever soil, precipitation and other climatic conditions do not forbid tree growth and where agriculture is sufficiently developed to need the protective influence of trees. All of the evidence available from studies that have been made and from the experiences of farmers and ranchers who have shelterbelts indicate that the plan for shelterbelts in the Great Plains is sound.

Specifications

Foresters think that trees should be planted close together so that the branches will grow together quickly and eliminate the necessity of cultivation. Foresters also claim that close planting with closed borders and complete protection from fire and grazing will develop a "forest" soil. Such a soil with its litter cover and permeable physical structure will conserve and store moisture to be used for the growth of the trees.

The spacing of early plantings included a variety of widths between rows, of spacing within rows, and of number of rows. Shelterbelts contained from 1 to 20 rows and these rows were from 8 to 12 feet apart with 10-foot isolation strips on each side. The distances apart in the rows of 2 to 8 feet were varied with the species. One, two and three row belts have been success-

refully
ing and
plant-
l.
ized in
legisla-
in an
In the
out 30
s civil-
y large
ts and
ic loss
plant-
re this
3, the
ropria-
riment
f 1924
plant-
ntings
were
wever,
to the
y set-
Plains
other
h and
need
e evi-
made
chers
n for

anted
w to-
f cul-
nting
from
Such
ysical
to be
a va-
ithin
con-
were
ation
the
ecies.
cess-



fully planted and maintained and have been effective in Oklahoma and Texas. Further north where snows are frequent, wider belts tend to collect the snow in drifts thus materially increasing the moisture for tree growth. The loss of a few trees in a narrow belt, which permits the wind to sweep through the openings, may destroy

Shelterbelts such as this help to harness the wild winds that come roaring across the plains. They protect land, wildlife and man, and at the same time add a touch of beauty.

A barrier against the fiercest wind, as reared in Burt County, Neb. The trees form a triangle-shaped shelter, with the slower-growing species at the left and the more rapidly-growing species at the right.



the effectiveness of the entire shelterbelt. The additional land and expense required to plant a few extra rows is the cost of insurance against possible complete or partial failure. Five to ten rows varying in height from shrubs to tall trees are now favored over the wider belts that were established a few years ago. The graduation in height from low shrubs on the outside of the belt to high trees in the center imparts an upward sweep in the air currents. The effect is felt to some extent at distances to leeward as much as 50 times the height of the trees. Twenty times, however, is usually considered to be the practical limit of effect.

We know now that zones or solid belts of trees from the North to the South as originally contemplated would have been a mistake, and in fact such planting never was attempted. The direction of the belts is carefully selected in relation to the prevailing winds. Sometimes they run North and South, sometimes East and West, and at many locations they are established in both directions to control winds during different periods of the year. The important problem is to locate the trees so that most of the air currents are dispersed before they get strong enough to damage agricultural crops. East-West belts frequently give this protection against summer winds but will not provide protection against the quartering winds of winter. It is because of the variableness of wind direction that belts in both directions are necessary. The frequency of belts depends upon the local conditions and the degree of protection necessary. Usually the intervals do not exceed 1/4 mile and under some conditions it may be advisable to have them as close together as 1/16 of a mile.

Species and Arrangement

In planning a shelterbelt, certain kinds of trees and shrubs are better than others. A great number of shrubs and evergreen and broadleaf trees have been planted during the past 50 years with varying degrees of success. The choice of species depends upon their adaptability to climate and soil, their susceptibility to insect and fungus damage, their economic and aesthetic values, and the degree of protection required. The hardwood trees found to be most useful in the Plains are cottonwood, green ash, American and Chinese elm and honey locust. Evergreens that have thrived best are eastern red cedar, Rocky Mountain red cedar, Austrian and ponderosa pine. Shrubs best adapted are Russian olive, Siberian pea, wild plum, lilac and choke cherry. The arrangement of rows us-

ually provide shrubs on the outside, evergreens next, tallest broadleaf trees in the center.

Protection and Care

Shelterbelts must be protected and, until well established, they should be cultivated. Trees will not live or a shelterbelt be effective if browsed and trampled by livestock. In early stages the trees are easily killed by competition of grass and weeds. Three to 5 careful cultivations annually are needed during the first 5 to 6 years after planting. This removes weed competition, aerates the soil, and puts it in better condition for holding moisture. Dead and diseased trees should be removed. The trees in the interior rows may be pruned, but no pruning should be done on the outside trees nor on any of the evergreens or shrubs. Trees that die during the early part of the life of the shelterbelt should be replaced the following planting season. When planted close together the trees crowd each other and suppressed trees will eventually die. Suppressed trees may be removed while the wood is still sound.

Influences

The direct and indirect results from shelterbelts in the Great Plains have been studied by numerous federal and local agencies. Many of these surveys pointed to benefits that were so great that most of the results were never published. Long time records of the influence of shelterbelts on crop yields in the United States are not available. The experiments that have been carried out indicate that shelterbelts not only hold the soil in place and control wind erosion but, in addition, crop yields have been increased, livestock has required less winter feed, farm buildings have needed less fuel, and farm gardens have produced more abundantly. One careful survey conducted in South Dakota shows that farmers who have shelterbelts believe that trees add \$1,000 to the value of each 160 acres.

Shelterbelts become a source of post and fuel material for the farmer after the belts have attained an age of 10 years and older. Desirable wildlife, game and insectivorous birds seek protection and propagate within the shelterbelts. This wildlife frequently is a source of food and recreation and the aesthetic value of the trees should not be discounted. In fact shelterbelts offer shade from the sun, shelter from the wind, improved appearance to the area, and increased value to the land. These factors add to the stability of farm life.

Agriculture in the Plains States without trees is hazardous. The best proof of the value of shelterbelts comes from the farmers who have them. These farmers even under war time restrictions and manpower shortage and without governmental aid except for free trees, are continuing to plant shelterbelt trees by the hundreds of thousands.

Disadvantages

Most of the disadvantages reported as a result of shelterbelts arise from badly selected locations. On any farm, trees in the wrong places are a handicap, not a help. It is true that shelterbelts break up large fields into tracts that are inconvenient and economically difficult to handle with heavy machinery. In some locations they may cause snow to collect in drifts on roads and about buildings, and may encourage late spring frost if they cut off air movements near the ground. However, the positive disadvantages of shelterbelts are so few and so trifling when compared with the benefits that only brief mention is necessary. In fact, the shelterbelts that have been established in the Great Plains stand as markers to the practical ideal of the pioneers who strove to bring forest benefits to a treeless region.

Future Plantings

Shelterbelts have been incorporated into the conservation program administered by soil conservation districts, with excellent results. The planting of shelterbelts as a separate project was discontinued in 1942. Since that time no special appropriations have been available but the work continues to go forward as a part of the farm conservation program that the Soil Conservation Service is carrying on by assisting soil conservation districts. Up to March 1, 1945, soil conservation districts have been organized over almost 45 percent of the area included in the Prairie States Forestry Project. New soil conservation districts are being formed rapidly within the remaining 55 percent of the area and indications now are that within a year and a half after the war between 60 and 80 percent of the Great Plains probably will be covered by locally governed soil conservation districts.

Outside of organized soil conservation districts the Soil Conservation Service is assisting farmers to establish shelterbelts to a limited extent as a part of the widespread conservation program. The Service has assisted 44,500 farmers and the Prairie States Forestry Project during its active life has helped 33,000 to establish shelterbelts.

Experience over a period of 10 years leads to the conclusion that a shelterbelt program sponsored and directed by local farmers and land owners with broad over-all federal coordination is more successful than a huge federal one-practice project. Programs sponsored by local groups under the leadership of the people who have a direct interest in the land are built upon a firm foundation. This has been proved by the Soil Conservation Districts in the Northern Great Plains during the war years of 1943 and 1944 where, notwithstanding the handicap of shortages of manpower and planting stock, more than 5,000,000 trees have been planted.


Shelterbelts are far from being a cure-all for agricultural problems in the Great Plains. They constitute but one of the many conservation practices that are being recommended. Other practices include rough tillage, crop rotations, cover crops, pasture contouring, terracing, diversion ditches, stock water tanks and ponds and many others. Correct land use, according to the needs and adaptability of the land, is essential on each farm. To give technical assistance to effect these practices is the foundation of the integrated farm conservation plan that is the basis of the work of the Soil Conservation Service, not only on the Great Plains, but throughout the United States.

Pinning Down the Sand

"The Dunes Stood Still" is the title of an article by Neil M. Clark in the April 28 issue of The Saturday Evening Post. It describes the levelling and successful regrassing of the Caddo dunes in Colorado which permitted operation of a railroad through the area and greatly simplified a problem in war transportation.

Mr. Clark writes: "Soil experts say we have made millions of acres of deserts in this country by overplowing and overgrazing low-rain-fall areas of the West. Lately the United States Soil Conservation Service gave dramatic proof that we can also unmake deserts. . . . The cost of unmaking this particular desert was \$131,250. Army engineers officially estimate that if they had had to use the alternative route, the cost would have been at least \$1,500,000 more."

Old West Lives Again in LU Projects



Working together, a group of livestock operators brand and vaccinate calves out on the open range. This is on one of the common-use pastures in one of the land utilization projects in the Northern Great Plains.

By O. E. McCONNELL

A tradition of the Old West was the use of rangelands in huge, unfenced areas. It may have been haphazard and uncontrolled, but the system had the advantage of flexibility. To some extent, this tradition of the Old West is being revived in the land utilization projects in the Northern Great Plains through the development of common grazing areas.

Common grazing is the term applied where two or more operators graze their livestock on the same range at the same time. Cattle belonging to several operators run at large and intermingle. Sheep are herded in bands by individual owners, or a band may be made up of several small flocks belonging to different people. Brands identify the individual ownerships. Under present land ownership patterns, not all units lend themselves to common grazing but to the extent that it is practical this plan is fostered in the land utilization projects.

Until the introduction of barbed wire fences common grazing was practiced as a necessity on the Public Domain lands before enactment of the Taylor Grazing Act. Frequently it led to bitter argument and combat between cattle and sheep operators and a consequent segregation of cattle and sheep ranges. Nevertheless, although cat-



Here's a well-bred youngster receiving the owner's brand and being vaccinated against disease.

tle and sheep are able to thrive together on the same range, sometimes in the past their owners have been unable to do so.

In those areas which were heavily homesteaded, the open or Public Domain lands that remained were in small scattered tracts and common grazing had almost disappeared. This trend has been still further accentuated in the last few years of abundant moisture, increased vegetation, increased numbers of livestock, higher prices for livestock and plentiful supplies of ready money, following the period of drought, low prices, abandonment of lands and a heavy relief load, which have witnessed the swing by a large number of operators in the Northern Great Plains toward individual grazing allotments and fenced-in units

EDITOR'S NOTE.—The author is chief, land management division, Soil Conservation Service, Lincoln, Nebr.

—contrary to the currently popular song, "Don't Fence Me In."

The years when the Federal government had to purchase a large percentage of the livestock and furnish relief of all kinds, with lands tax delinquent in several states and going back to county ownership in wholesale lots, are still remembered by some of the people. And a considerable blocking in of units by the purchase of county and other lands is taking place. Ranchers are taking advantage of the opportunity to buy the land they need while they have the money to pay for it.

Whether common grazing or the individual-unit system is most economical in the long run seems to be of secondary importance to those men. Perhaps those ideas born of our way of life, which makes us Americans and in turn makes the United States what it is—that indescribable feeling we

call independence, freedom, "mine to do with as I please"—has more to do with the individual's desire for a little domain he can call his own without undue outside interference.

The land utilization projects, however, are also furnishing the opportunity for demonstrating that people can still work together and gain the advantages of cooperation which is also a part of our way of life.

The Soil Conservation Service is custodian for 4,600,000 acres of lands in North and South Dakota, Nebraska, Kansas, Wyoming and Montana, which were purchased by the Federal government under Title III of the Bankhead-Jones Farm Tenant Act. These lands are the nucleus for 18 land utilization projects totaling 20,000,000 acres and embracing also privately-owned, county, Public Domain, and state lands. They are principally grazing and commensurate feed-base lands.



Roundup on one of the common-use pastures, preparatory to branding on open range.

Practically all of these lands, including the government-owned acres, are administered by 44 local governing bodies—grazing districts, soil conservation districts or grazing associations, organized under state law. The 156 common grazing areas which have been set up total 5,000,000 acres—1,600,000 acres of which are Title III lands. The areas vary in size from 2,000 to 70,000 acres, each accommodating the livestock of from 2 to 28 operators which totals 100,000 cattle, 150,000 sheep and 6,000 horses. Some of the grazing districts have no common grazing; in others, nearly all of the summer grazing lands are used in common. Many of these smaller operators are located so that the pasture needed for summer grazing is not available close to the headquarters. Where this is true, the common grazing area is the solution of their difficulties—a place to put their stock where it will be properly cared for, even though quite far from the headquarters.

Developments on the lands within the common grazing areas have been carried on cooperatively by the districts and Government. These include the construction of corrals in many of the common grazing areas, dams and wells and some dipping vats. It is planned to have corrals in all of the larger common grazing areas to facilitate branding, vaccinating, rounding up for shipment, and so forth. For some of the larger areas, the districts employ range riders to look after livestock and maintain the improvements. Extra district fees, above those charged by the government for the forage, are charged for the additional service where salt is furnished, riders employed and improvements maintained. All this work is necessary for handling livestock, whether it be done individually or in groups.

Increasing the quality of stock produced on the common grazing areas is centered in the bull committees provided for in each district's by-laws. The committee is either elected or appointed by the district board of directors, and usually one of its members is the county agent or some other person trained in agriculture. The other members are experienced operators. The duty of the committee is to pass upon the quality and conformation of all bulls turned into the common grazing areas. The use of inferior bulls is discouraged or not permitted in these pastures.

In some districts bulls must be registered animals of the breed decided upon by the majority of the operators using each pasture. While bulls owned by some of the better operators are of higher quality, those generally in use now in com-

mon grazing areas are average quality or better and the quality of the cattle as a whole has been increased materially. Of course, not all bull committees have made the same progress. Some are doing an excellent job; others are still somewhat lax. The present higher prices of better bulls has retarded all the progress desired in this direction. An operator, however, may use an inferior bull so long as he keeps it within the confines of his own fenced unit.

Common grazing saves much, perhaps an average of 50 percent, in fence construction and maintenance. The extent of the saving depends on the size of the common grazing area and the number of operators it accommodates. Savings in water developments are about the same. Since livestock water is one of the important limiting factors governing use of the range in the Northern Great Plains, it deserves careful consideration. In addition to the savings in the construction and maintenance of fences and reduction in the necessary number of water developments, compared with individual fenced units, common grazing permits taking advantage of better sites for water developments. Thus, better located and more dependable livestock water supplies result.

The common grazing system has some district advantages in addition to cost savings for the smaller operator. It has a tendency to get people and communities in the habit of working together, which makes for a better community in which to live. It also leads to the adoption of one breed of livestock in a community. There are several good breeds, but standardizing on one breed in a community is nearly always better than a helter-skelter intermingling of two or more breeds.

Further, grazing is better controlled with respect to season of use and degree of stocking in the common grazing areas than where the land is divided into individual fenced-in units. This better management increases the chances for continued forage production at a high level.

Last, but by no means least, the use of common grazing areas acts to protect the winter feed supplies. Usually the common grazing areas are located some distance from the headquarters, so that the stock are out of the way during the summer grazing season, which is also the growing season for feed crops. Operators, then, are free to go about the business of raising the feed needed to winter the livestock. Likewise, pasture reserved for winter use is not grazed, and is waiting for the stock when they are brought back from summer range. Too often, in individual units, pasture



The branding irons of 10 different owners are in this fire.

which should be reserved for winter is used in summer. This creates a scarcity of winter feed. The result, in many cases, is winter range used beyond the safety point and damaged, and livestock in poor condition.

Thus, through cooperation, the operator of a livestock unit is enjoying some of the tradition

of the old West through cooperation with others in land utilization projects. He is, at the same time, gaining the protection arising from locally controlled use of the grasslands. And he is pointing a way for other communities who may wish to solve similar land used problems.



Purebred bulls acquired by grazing association in effort to improve quality of livestock in area.



Corrals in which to separate animals for branding, vaccinating and dehorning, and for cutting out stock to be shipped, are a regular feature of the common-use pastures on land utilization projects. Here cattle for shipping are being separated from the foundation herd.



FIGHTING FERN WITHOUT FIRE

The fern-covered areas are a constant fire hazard until converted to productive use. Some lands can be cropped or put into permanent pasture. Slopes like that in the background are often suitable for timber.

By ALBERT ARNST

In the Douglas fir region of the Pacific Northwest grows a plant that is a blessing to foresters and a curse to farmers.

It is fern (*Pteridium aquilinum* var. *pubescens*)—also called bracken, brake fern, western bracken and hay brake. It has no commercial value; but the successful growth of Douglas fir, western hemlock, spruce and western red cedar seedlings depends upon its shade and protection in many areas. The problem, then, is how to eradicate this troublesome plant from farmland without resorting to fire that spreads into adjacent seedling forest areas and destroys tomorrow's timber crop.

Bracken fern is an insidious and persistent resident of cutover lands and forest intermingled rural areas. It is found anywhere and everywhere in that region, chiefly west of the Cascade Mountains, and grows luxuriantly in the Coast Mountains of Oregon. It clings to farmland and pasture land with a tenacity that matches the fury of the farmer who seeks to stamp it out, and it clothes

logged and burned forest lands with a dense cover which frequently tops a man's height. Tall tales are told of farmers' livestock lost in fern fields and of foresters' cruising the swaying stalks, but even Paul Bunyan agreed that the plant reaches amazing heights.

The plant reproduces by means of creeping root stocks buried deep in the ground and also by spores carried in the characteristic small brown pustules on the underside of the fronds. In its proper place on forest adapted lands fern is a valuable nurse crop for tender seedlings getting started on recently logged and slash burned areas. Extended studies by foresters have shown that fern shade nurtures more trees and bigger trees. Once the trees get above the fern canopy, they eventually shade it out and finally eliminate it from the timber stand.

But fern is a menace to land use managers, particularly in soil conservation districts, because in winter and early spring it becomes a serious fire hazard. The tall stalks die out in fall and eventually are broken down by wind and rain. Crumpled to the ground in densely interlaced masses, they dry out rapidly under changing humidity and

EDITOR'S NOTE.—The author is forestry specialist, Soil Conservation Service, Sedro-Woolley, Wash.

weather conditions. When tinder dry, they constitute a dreaded "flash fuel," which, when touched off by a spark, will carry fire with utmost swiftness.

In the fire's wake may be found the thousands of little scorched trees that, unseen to the casual glance, were snuggling under the fern for protection. Fern fires also consume larger trees and destroy much wildlife. During deer fawning season, fires kill fawns, or perhaps nesting game birds, such as grouse, quail and Chinese and native pheasants. Burning fern destroys ground cover which has food value for wildlife, and it also muddies clear streams with erosion runoff. No one profits when a fern fire races across a distant slope.

Of additional interest to farmers is the fact that fern, growing or dried, is considered poisonous to horses, sheep and cattle. Sunlight seems to aggravate the poisoning effect, which results in loss of control of the leg muscles, constipation and eye pupil dilation, but causes no fever. Fern, there-

fore, has no place in conservation farming, because its presence on agriculturally adapted lands is a sign of improper land use.

So-called burning weather in February and March makes match-happy ranchers squint at fern fields with fire in their eye. Soil conservation districts in fern areas therefore must face fern-burning realistically, for some ranchers are potential "fernomaniacs." Statistics show that most fern fires start primarily on agricultural or assumed agricultural lands, and through carelessness, spread to nearby fields of fern-protected tree seedlings.

John Farmer generally burns fern, not to get rid of trees, but to eradicate the fern, which bothers his farming. Fire is his tool, because no one has given him a better weapon. To eliminate fern fires, the rancher must be shown a superior way to do something he will do anyway; and, furthermore, he must be convinced that on some land it is not sensible to fight fern with fire or any other method. Both the rancher's and the public's



Ferns actually nurse young seedlings like this one to the point where they can fight their own way up in the world —if they are lucky enough to escape fire.



Fire kills trees but not ferns. Lustier than ever, the ferns come back after a hot ground fire has destroyed all other growth and the surface layer of humus and duff.

appreciation of land values needs to be improved.

Each year in the Douglas fir region of the northwest, the Keep Oregon and Keep Washington Green Committees, through their cooperating agencies, carry on an active "Don't Burn Fern" campaign by press, radio and poster. A still more positive approach, equally convincing to farmer and forester, is demonstrating that fern can be banished from pasture or crop land more effectively by not using fire, and that better tree crops will mature sooner if fire is eliminated.

In soil conservation districts, soil inventories and land use capability classifications are front line weapons in the skirmish with fern burners. These indices rate the best long-time productive capacity of each farm field as determined by soil, erosion, slope and climate. They also show where crops can be grown successfully and where permanent vegetation, like trees or grasses, should replace row crops. This down-to-earth information proves it is not reasonable to develop remote "pasture" on fern covered timber growing soils, if better soiled fern land nearer the farmstead can be improved permanently with far less effort.

Fern can be licked by not using fire. On soils

adapted for pasture and on tillable slopes, plow the land deeply and then harrow out, pile up and burn the rootstocks. Cultivate the land frequently during the remainder of the season or grow a row crop. Effective anti-fern treatment is the growing of oats and vetch, cut early for hay, followed by summer fallow.

Mow and clip slopes that are not too steep for handling in this manner. Run the mower over the green fern patch around July fourth and again later in the season. This bleeds the plant and retards its recovery, because the bracken is hit while manufacturing food for its buried rootstocks.

Fern burning may be sanctioned as a one-time treatment on still steeper land, if it is adapted for permanent pasture and if the rancher will follow the fire warden's advice. Burn in the fall, seed immediately to permanent grasses and pasture them when they become established. Fern-hating sod formers are used—Chewings fescue, alta fescue, creeping red fescue, highland bentgrass, orchardgrass and tall oatgrass. Annual grasses have no value for fern control.

These fern eradication techniques save trees. The field formerly torched may not have had little trees in it, but probably it bordered on a steeper slope bristling with fern-sheltered seedlings. Fire kept out of the first field also would have been kept out of the second.

Tree farmers are shown hundreds of seedlings snuggling under the dried fern. They learn that repeated fires kill all trees, impoverish soil and stimulate fern growth, because the rootstocks escape injury. They appreciate unburned fern as a nurse crop in growing timber on forest adapted lands. They know that tomorrow's timber crop may be found under today's fern patch, and that some day some community and some industry is going to benefit from the harvest of the matured trees.

Europe's Needs Reported

"The needs of Northwest Europe's liberated areas are grave, not only from a humanitarian aspect, but because they involve internal and international political considerations. The future permanent peace of Europe depends largely upon restoration of the economy of these countries, including a reasonable standard of living and employment. United States economy, too, will be deeply affected unless Northwest Europe again resumes its place in the international exchange of goods and services.—From report of Judge Samuel I. Rosenman to President Truman.

Pea
plasti
Cross
when
with
farmer
The
pletin
classe
blowing
War
crease
soil co
sulted
maint
Con
soil is
Timbe
rotati
croppi
impro
struct
proble
W.
Leon
sistan
he ha

EDITOR
Service,

GROWING PEANUTS THE CONSERVATION WAY



This field is stripped with eight rows of peanuts and four of legumes, of which the two rows outside are crotalaria and the two inside are brabham cowpeas.

By W. M. NIXON

Peanuts, used in products ranging from soap to plastics and anti-freeze, grow best in the West Cross Timbers sector of Texas and Oklahoma when they are produced the conservation way—with a complete soil conservation program, farmers in the area have found.

The problem of producing peanuts without depleting the land of this area, which already classed as critical from the standpoint of soil blowing and washing, faced farmers when the War Food Administration first asked for increased production. The application of a complete soil conservation program for each farm has resulted in increased production of peanuts while maintaining and improving soil fertility.

Combating the potential blow-off of fertile topsoil is one of the chief problems in the West Cross Timbers. Farmers here rely on a program of crop rotation, proper management of crop residue, strip cropping, contour farming, and cover and soil-improving crops to prevent loss. Terraces are constructed on those areas where water erosion is a problem.

W. A. Maples is a cooperator with the Upper Leon Soil Conservation District. With the assistance of Soil Conservation Service technicians, he has established a complete soil conservation

program on his farm, 10 miles northeast of Rising Star, Tex. This program paid off to the extent of a 70 percent increase in peanuts and an 85 percent increase in hay production in 1944.

Mr. Maples produced an average of 44 bushels of peanuts per acre in 1944 following crotalaria strips. He obtained 30 bushels per acre where peanuts followed peanuts.

Some of Mr. Maples' practices to secure better harvests include interplanting cowpeas and sorghum in strips alternating with strips of peanuts. The sorghum and pea strips afford protection against wind erosion. The peas supply nitrogen which keeps the sorghum from having a detrimental effect upon the following crop. He has used phosphate on winter cover crops of rye mixed with hairy vetch or Austrian winter peas. The cover crop was broadcast ahead of peanut digging. The hairy vetch and Austrian winter peas produce nitrogen, one of the most expensive fertilizer elements, now scarce because of its extensive use in manufacture of explosives. Mr. Maples has practiced late land preparation on deep blow sand, and he cuts his strips of sorghum stalks so they remain near the surface of the soil, forming a trashy mulch which conserves moisture and gives protection from hard rains and high winds. His experiences are typical of this area, where sandy soil, subject to severe wind erosion, predominates.

H. H. Lawson, cooperator with the Brown-Mills Soil Conservation District near Rising Star, Tex.,

EDITOR'S NOTE.—The author is regional agronomist, Soil Conservation Service, Fort Worth, Tex.



Typical of many fields of the West Cross Timbers area, where conservation measures are not used on peanut lands. After harvesting the fields stand bare and topsoil drifts to fence rows. A faint wash in the center of the field, which had been solidly in peanuts for two consecutive years, indicates both water and wind erosion.

uses a cover and soil-improving crop of rye and vetch to prevent blowing and washing. Organic matter and nitrogen are added to the soil when the crop of rye and vetch is turned under as green manure.

Mr. Lawson got 26 bushels of peanuts and 880 pounds of hay per acre from peanuts following rye alone. The peanuts following fertilized rye and vetch produced 44 bushels per acre and 1,628 pounds of hay. While the yield from the peanuts following rye alone produced at an above-average rate, the crop Mr. Lawson harvested following addition of vetch and application of fertilizer showed an appreciable increase.

An effective strip-cropping system for the Texas-Oklahoma Cross Timbers section consists of 4 to 8 rows of peanuts and 4 of corn, sorghum or crotalaria.

Grain sorghum grown on the contour in strips

of 4 rows alternating with 8 of peanuts gives good protection against wind erosion. A rotation is provided by moving the strips each year.

The farm of C. M. Caraway, Sr., 6 miles east of DeLeon, Tex., provides a good "before and after" example of the benefits of peanut grown the soil conservation way. Mr. Caraway is a cooperator with the Upper Leon Soil Conservation District and also is a member of the State Soil Conservation Board.

On land planted continuously to peanuts for 10 years, he harvested 7 to 8 bushels of peanuts and 10 bales of hay per acre. On a field where the soils and slopes were similar but where such practices as contour cultivation, cover and soil-improving crops, and crop rotations have been in use about 20 years, Mr. Caraway got a yield of 30 bushels of peanuts and 30 bales of peanut hay per acre.

IMPROVE GRASSLAND FOR BETTER LIVESTOCK

By R. H. LUSH

The necessity for providing sufficient food for civilians, for our armed forces, and, in so far as possible, for our Allies and the liberated peoples is self-evident. It gives new emphasis to the desirability of feeding the soil to improve grasslands. A well-fed soil produces strong plants which, in turn, contribute largely to the health of livestock.

EDITOR'S NOTE.—The author is pasture specialist, The National Fertilizer Association, Washington, D. C.

Completing the cycle, livestock and livestock products normally furnish a large proportion of the energy, the protein, the phosphorous and particularly the calcium in the average diet.

More livestock products are needed, but there is a shortage of hay, protein feed and labor with which to make the greatest production under normal methods of feeding. Of the total feed consumed by animals during the 3-year period of 1938-40, hay, pasture and other forage crops supplied approximately 60 percent. Emphasis should

be giv
espec
system
hogs
feed a
grazin
Ex
of the
able.
the f
than
pecial
exper
hay c
fertil
the in
Of
slight
7 per
more
about
State
acres
impro
or bor
three
65 per
1.2 ac
cludin
there
Mill
rainfa
be eco
in the
presen
tenanc
age is
too ser
Soil co
million
the Un
able an
another
improve
croplan
or to g
load of
acres.
More
crop ac
areas,
as well
grain a
able fo

be given to increasing this part of the feed supply, especially for cattle and sheep, and management systems devised which would make it possible for hogs and poultry to obtain 25 percent of their feed and nearly 50 percent of their protein from grazing crops.

Experimental results are available in nearly all of the States to show this to be entirely practicable. Furthermore, the cost of producing feed in the form of pasture and hay is considerably less than the cost of producing grain, and this is especially true of the labor cost. Results of many experiments indicate that yields of pasture and hay can be increased greatly by the proper use of fertilizer and lime where it is needed, and that the increases obtained are profitable.

Of the total land area of the United States, slightly more than half is in grass, of which only 7 percent is in plowable farm pasture, slightly more than that in farm woodland pasture, and about 20 percent in other pasture. The rest is State and Federal grazing land. Of the 130 million acres of plowable pasture land most likely to be improved, 80 million are in the 31 States east of, or bordering, the Mississippi River, and in the three Pacific Coast States. These States include 65 percent of the Nation's livestock but have only 1.2 acres of plowable pasture per animal unit. Including the 100 million acres of other pasture there are only 2.7 acres per animal unit.

Millions of these acres are in areas of variable rainfall where the addition of fertilizer might not be economical in some seasons. Other millions are in the Middle West, where lack of soil fertility is presently becoming a serious problem in the maintenance of high yields. Much of the pasture acreage is abandoned cropland too poorly drained or too seriously eroded to be made quickly productive. Soil conservation surveys show that, although 398 million acres are available for crop production in the United States, 40 million of these are unsuitable and should be retired to grass and woodland; another 350 million acres of land are subject to impoverishing erosion. Abandoned and eroding cropland needs to be retired to perennial grasses or to grasses and legumes. This throws a heavier load of production on remaining crop and pasture acres.

More efficient use should be made of the present crop acreage, along with conservation of "waste" areas. Farmers need range animals of the West as well as home-raised livestock, to convert some grain and forage into highly nutritious and palatable food, to help maintain soil fertility, to dis-

tribute labor, save transportation, and to add to income. The use of pasture saves labor and protein feed, supplies adequate quantities of known vitamins and most minerals where the soil is fertile or fertilized, aids in sanitation, conserves soil and farm machinery. Moreover, pasture is usually available when needed most and is the cheapest feed for all classes of livestock.

Improvement of pastures and hayland by use of fertilizer and lime has progressed rapidly during recent years. In 1929, less than 200,000 tons of fertilizer were used on about 2 million acres of pasture and hayland; in 1942, nearly 1,300,000 tons were used on about 11,000,000 acres; 1,500,000 tons in 1944. But this is only a start. A post-war planning committee of the U. S. Department of Agriculture has suggested that nearly one-half of the total commercial plant food should be applied to grassland and other close-growing crops. The National Planning Association reports that 5 times as much fertilizer could be effectively used on pastures and haylands as in 1942.

About one-twelfth of the combined plowable pasture and hayland acreage is now being fertilized. There are millions of acres of non-plowable pasture that might be fertilized profitably. Still other millions of acres of supplementary pasture and hay crops are fertilized but little, if at all.

A compilation by the War Food Administration of many experiments with fertilizer on pasture and hay crops shows that each pound of plant food used in the form of complete fertilizer brought about an increase of from 11 to 22 pounds of dry matter and 1 to 3 pounds of digestible protein. Large increases were obtained in the Northeastern and Southern States. The chief response in the Western States was to the use of phosphoric acid. There were wide variations between States and between experiments. Each pound of nitrogen, of which there will be a large supply in the future, made possible from 14 to 20 pounds of digestible nutrients, containing 1.8 to 3.1 pounds of digestible protein—a very profitable exchange for feed.

With the high cost of labor and of legume seed, it is now desirable to prolong old legume stands by fertilizing or by reseedling with an adapted grass mixture, such as brome grass, orchard grass, timothy, or ryegrass. This will give additional hay and grazing, with a somewhat longer rotation. In postwar farming it may be that the quickest and cheapest way of obtaining additional feed will be to apply liberal quantities of complete fertilizer to new and old sod crops, supplemented by nitrogen top dressing. The fertilizer not only in-

creases the yield but also insures that better soil cover which aids in erosion control.

More than 40 million acres of small grain are seeded annually in the States where there is sufficient rainfall to insure profitable results from fertilizer use. It is estimated that fully 5 million of these acres are seeded primarily for grazing or hay or soil improvement. Fertilizer applied at seeding time insures a better stand, an earlier fall cover and consequent soil protection, fall grazing and, in the South, winter grazing. In general, there is less winter killing. If animals are kept off the fields in wet weather, some winter and early spring grazing can be obtained without reduction of grain yield or injury to new seedings. Feed thus obtained is cheap. It supplements permanent pasture and roughage feeding, aids in manure distribution, and for dairy cows insures better quality of milk.

Animals thrive according to the extent to which the soil supplies the essential elements to grains utilized as feed. Past emphasis has been given to the development of vegetation by crop species and tonnage yields, rather than by nutritive values. Yet, poor soils make poor feeds and unprofitable livestock. A depletion of fertility in pastures is constantly under way. Each pound of meat, each quart of milk, removes plant food from the soil.

Better nutrition should start at the grass roots, with deficiencies made up by increasing the supply of nutrients. At Joliet, Ill., a ton of bluegrass hay from a plot treated in early spring with complete fertilizer contained 435 pounds of protein and 6.4 pounds of phosphorus, as compared with 205 pounds of protein and 3.6 pounds of phosphorus per ton of hay from untreated land.

When soil treatments were made in Missouri on adjoining plots, their effects were registered in sheep as differences in animal growth per unit of feed consumed and as variations in the quality of the wool. At the Georgia Experiment Station beef cows on permanent clover pasture treated with limestone and phosphate produced a 100 percent calf crop two years in succession as compared with only 50 percent of a calf crop for cows on unfertilized pasture. Almost as striking differences occurred on fertilized and untreated ranges of the King Ranch in Texas. A blind horse in Ohio grazed only to the edge of the fertilized pasture area and avoided the untreated grass. Animals have "horse sense" about feeding, but farmers have only recently become aware of the differences in feed due to soil fertility. The addition of plant food to

the soil where it is necessary might be termed as livestock health insurance, as well as profitable practice.

Grain must, of course, be fed to fattening hogs and poultry, heavy-producing dairy cows, some work stock, and livestock not on farms. All other livestock can use a larger proportion of forage crops, if every effort is made to give more days of good grazing on each pasture acre, and to harvest a higher quality and yield of hay.

WATER STORAGE IN THE SURFACE SOIL

By O. R. NEAL

The matter of soil-water relationships has been the subject of extended investigations on the part of soil physicists for many years past. In Bauer's *Soil Physics* investigations on physical properties of the soil in general and on moisture relationships in particular are traced from the work of Davy in 1813 and Shubler in 1833 up to the present time. The first extensive studies on physical properties of the soil were made by Wollny between 1879 and 1898.

Wollny studied the properties of soils with a view to the effect on plants. He was able to present a reasonably accurate picture of the importance of soils and plants on the disposition of rainfall or, in modern terminology, the role of soils and plants in the hydrologic cycle. He measured the interception of rainfall by vegetation and studied the protective effect of vegetative canopy on surface soil porosity. He measured the quantities of runoff and erosion from plots of different degrees and direction of slope, with different soil and crop conditions. Lysimeters were used to study the movement of water through soil under different crops and treatments. A surprising number of recent contributions in the fields of hydrology and conservation are actually only confirmations of Wollny's original work.

During the period of Wollny's work in Germany several American investigators were concerned to a greater or less extent with soil moisture studies. The best known of these were Hilgard of California and King of Wisconsin. King is generally recognized as the father of soil physics in this country. His book entitled *Physics of Agricul-*

EDITOR'S NOTE.—The author is project supervisor, Soil Conservation Service, New Brunswick, N. J.

ture enjoyed widespread popularity and is of interest at present, particularly in connection with the author's view on soil moisture and tillage problems.

Beginning at about 1900 there was a marked decrease in research and presumably in interest in physical properties of soils. From that date up to about ten or fifteen years ago, most of the soil physics work in this country was done in the Bureau of Soils. Briggs, McLane, Shantz, Buckingham and others made notable contributions during this time, particularly in the field of soil moisture properties. Briggs and his associates developed the moisture-equivalent technique as a measure of the ability of soils to hold water. They also introduced the concept of the wilting coefficient to express the moisture content of the soil at which plants permanently wilt. Veihmeyer and Hendrickson suggested the term field capacity to designate the moisture content of the soil after the rate of downward movement of added water had decreased to a very low value.

In any practical consideration of soil moisture in relation to crop growth, whether the moisture is supplied by natural precipitation or by irrigation, the field-capacity and wilting-point values are of considerable significance since they mark the upper and lower limits of soil moisture usable by plants. In any particular soil each of these values is more or less fixed as a result of the physical nature of the soil. Treatments which tend to increase the field capacity usually result in a more or less equivalent increase of the wilting point. As a result, the quantity of water that the soil can hold for use by plants is not materially changed.

When water is added to the surface soil, it moves downward only as each successive layer of soil is raised in moisture content to the field capacity. It is not possible to raise the moisture content of the surface foot, for example, by a small amount and have the added moisture distributed uniformly through the layer. Rather the added water will all be held in the top one or two or three inches depending on the amount added. In a soil weighing 90 pounds per cubic foot and having a field capacity of 35 percent, approximately 6 inches of water can be held in a surface foot of soil. If the soil were completely dry, as is rarely if ever the case, an application of 2 surface inches of water would all be held in the first four inches of soil. Under actual field conditions the moisture content at the time of water addition would be somewhere above the wilting point, with the result that from 2 to 3 inches of water would be present

THE GIST OF IT

This article points out that certain moisture constants of the soil are of particular importance in connection with irrigation. The two most important are field-capacity and wilting-point. The field-capacity, for all practical purposes, represents the only moisture content that can be induced in field soils by the addition of water. The wilting-point represents the condition below which plants are unable to obtain water at a rate sufficient to support growth processes. The range between the two values represents the quantity of water that the soil can hold for use by plants.

It is observed that water losses by evaporation are limited largely to moisture held in the surface layer. Attention is directed to this fact in the planning of irrigation applications.

Also noted are the effects of irrigation practices on physical properties of the soil, organic matter content, and general soil productivity. The need for adoption of effective conservation practices in connection with the use of supplemental irrigation is emphasized.

when the addition was made. Under these conditions 2 inches of added water would wet the soil thoroughly to a depth of about 8 inches.

Losses of moisture by evaporation from the soil surface occur largely at the expense of the water held in the surface 6 to 8 inches of soil. There may be a limited amount of water lost by evaporation from greater depths but the quantity is not of great practical significance. Losses from the surface layers, however, may be high. There is experimental evidence that losses of from 0.20 to 0.25 inch per day are not uncommon when the immediate surface of the soil is wet. In a lysimeter experiment conducted several years ago two soils of quite different profile characteristics were compared in respect to quantities of water lost by evaporation. A soil that was uniformly permeable throughout the column lost an annual average of 54 percent of the precipitation as surface evaporation.

The second soil which had a relatively impermeable layer at a depth of about 9 inches lost 71 percent of the precipitation. Presumably the impermeable layer tended to hold more of the water near the surface where it was subject to evaporation. The classical experiment in this connection

was reported by Veihmeyer a few years ago. He filled containers having a depth of about 30 inches with moist soil, then exposed them to evaporation but protected them from rainfall for periods varying from 80 to 167 days. At the expiration of such periods the dry soil on the immediate surface was mixed with the underlying soil and a crop of vetch was grown without the addition of water. These results indicate clearly that evaporation losses are limited to water held in the immediate surface layers.

In agricultural operations where rainfall alone is depended upon as a source of water the foregoing information concerning water-holding capacity of the soil and the losses by evaporation are of academic interest but have no direct field usefulness since the amount and time of water application cannot be controlled. *In the case of supplemental irrigation, however, the situation is entirely different.* The rate and quantity of water application can be manipulated in whatever fashion produces the greatest economic return. In connection with evaporation losses, for example, it has been shown that water is lost in appreciable quantities from only the surface few inches of the soil regardless of the total depth of water penetration. In a particular soil a 1-inch application of water might be held entirely in the surface 5 inches where it would be subject to rapid evaporation loss. Under the hot, drying conditions that often obtain during periods when irrigation is needed, it is conceivable that a large fraction of the added water would be lost in the course of a week. When another irrigation was made the process would be repeated. Only a small fraction of the added water would be available to plants under this condition. If, on the other hand, 2 inches of water was applied at the time of the first application the depth of penetration might be 10 to 12 inches. Surface evaporation would still be limited to the surface 5 inches, more or less, and all the water below that depth would remain for plant use. The efficiency of the operation from the standpoint of quantity of water actually made available for plant use would be considerably more than doubled by this procedure. The foregoing statements concern only the efficiency of water use and do not take into account the use of labor, quantity of pipe required, rate at which water can be obtained, and other factors.

Research information is sorely needed and, to my knowledge, is not available on many of the factors involved in the practice of supplemental irrigation. The rate at which water can be applied safely to different soils under different condi-

tions of cover is not known. Presumably it is desirable from the standpoint of the operator that water be applied as rapidly as possible without producing runoff. Time required for application of a unit quantity of water would be greater with heavy applications than with light ones. When water is first applied to a cultivated surface it is absorbed very rapidly, but the rate of entry decreases as the period of application is lengthened. In the case cited above concerning the application of 2 inches of water instead of 1 inch, the time required would doubtless be considerably more than doubled. Contouring, terracing, mulching, and other conservation practices will increase the rate of water penetration under irrigation just as they do for natural precipitation.

Information is needed on the quantity of water that can be held in the root zone of the crops that are being grown. This might vary considerably with different soils. Simple, rapid, and moderately accurate methods are needed for determining the moisture content of soils in the field in order to determine irrigation needs. Some progress has been made along this line. Plaster-of-paris blocks which are imbedded in the soil in a suitable container and can be removed for weighing offer some promise in this connection. Tensiometers are very useful for certain moisture levels but do not operate over the entire range of soil moisture from field capacity down to the wilting point. Other equipment and methods have been proposed for the measurement of soil moisture content and have been used with more or less success. There remains a need, however, for a simple field method of determining the moisture condition in the soil with respect to crop needs. Such a method, to be ideal, will be based on the newer energy-concept of moisture measurement rather than on a mere determination of weight of water per unit of soil.

In considering the developments that may be of service in connection with irrigation activities, the possible value of long-range weather forecasting should not be overlooked. We are told that great progress has been made in this science during recent years. It is not too fanciful to envisage the use of these techniques in agricultural operations. Thornthwaite and his associates have recently suggested a procedure by which deficiencies of soil moisture can be determined. In the humid areas where relatively short drought periods occur at irregular intervals it would be of great benefit to the farmers if such occurrences could be predicted with even moderate accuracy. Furthermore, accurate forecasting for short periods of only a week or more would serve as a guide for irrigation.

In any discussion of irrigation problems the need for conservation practices should be strongly emphasized. It has been shown repeatedly that the deterioration of soil structure and the exhaustion of organic matter are responsible, in large part, for increased erosion on our crop lands. The very nature of the irrigation practices is such that these deteriorative processes will be accelerated. In our efforts to encourage the maintenance of a good supply of organic matter in the soil, we sometimes lose sight of the fact that the decomposition of organic matter is a normal and desirable process. The following is quoted from an earlier statement on this subject:

"Organic-matter decomposition, with the resulting liberation of plant nutrients, is desirable and necessary under average conditions in order to maintain the soil in a well-aggregated condition which is resistant to erosion and in order to maintain a productive condition. The accumulation of a large amount of organic material that was highly resistant to decomposition would not necessarily contribute to good physical condition of the soil and to soil fertility. The desirable condition is one where a system of crop and soil management is so arranged that additions of good quality organic materials are made at regular intervals. The

normal decomposition processes that contribute to good physical condition of the soil and to high productivity can then proceed without danger of the eventual exhaustion of soil organic supply."

The need for soil management practices which will maintain organic matter is increased when irrigation is practiced. The maintenance of a favorable level of soil moisture during periods that, without irrigation, would be characterized by drought tends to increase the rate of organic matter decomposition.

Contouring, terracing and other mechanical conservation practices are highly desirable to prevent runoff both from irrigation water and from precipitation. Damage from runoff and erosion on some irrigated areas is serious. The quantity of irrigation water needed and consequently the expense of application is reduced as measures for more effective conservation of natural precipitation are adopted. It is conceivable that the apparent need for irrigation on many areas would be completely eliminated by the adoption of effective practices for the conservation of natural precipitation. Certainly, in view of both the immediate and the long-time benefits involved, adequate conservation practices should accompany the irrigation process.

ACADEMICIAN T. LYSENKO ON PLANTING OF WINTER WHEAT AND RYE IN UNPLOWED STUBBLE

Excerpts from "What is the Essence of Our Proposal Regarding Planting Winter Crops in Stubble in the Steppes of Siberia"

By Acad. T. Lysenko, pp. 25-36, *Sotzialisticheskoe Khoziaistvo*, Vol. 5-6, 1944

Translated and Abstracted
By D. B. KRIMGOLD

Subject of lively discussion among soil conservationists of late has been the question "To plow or not to plow" Faulkner's debatable book, "Plowman's Folly," has released a flood of discussion. The pros and cons have flown thick and fast. This book has served to focus attention on a fundamental operation of farming and has spurred an interest in a topic long due more general attention. Research scientists of the Soil Conservation Service have for a number of years been investigating various methods of planting crops without sub-

jecting them to unnecessary dangers of erosion. They have developed the so-called stubble mulch farming and have worked out a large number of variations of the "trashy cover" idea for use under a wide range of conditions.

Conservationists generally, and especially workers in the winter wheat regions of the United States, will therefore be interested in the discussion of a related subject by Academician Trofim Lysenko whom a national magazine includes among "some of the world's great contemporary scientists."

The first interesting thing about Lysenko's article is that it shows that he, the great Lysenko, is definitely on the defensive and must overcome the objections and criticisms of lesser lights. He begins his discussion thus:

"The question of planting winter crops on unplowed stubble in the prairie regions and the open sections of

EDITOR'S NOTE.—The translator is soil conservationist (research), Soil Conservation Service, Washington, D. C. The magazine "Sotzialisticheskoe Khoziaistvo" is the official organ of the Narkomzem (Department of Agriculture of the U. S. S. R.).

Siberia, Transural, and northern Kazakstan is one of national economic significance. Yet this problem faces a negative and often hostile attitude on the part of a number of agricultural workers and scientists and of agricultural agencies. This affects in certain measures the yields of cereals in these regions.

"Without entering into the essence of the difference between my proposal and the old well known method, I shall enumerate the basic reasons which are commonly given for low yields of crops planted on unplowed stubbles. These reasons are:

"1. During dry summers it is hard to obtain germination of winter crops on poorly plowed fallow. It therefore follows that for crops planted in stubble, (that is, in fields which are not worked at all after the harvest) there would certainly be poor germination. On such plantings germination can be obtained only in years with ample rainfall during the latter half of summer.

"2. Winter wheat in the prairie regions of Siberia seldom winters well even on well worked and early plowed fallow. Rye comes through well on 'black' early fallow but on late fallow, which is plowed immediately prior to planting, even rye does not winter well. Therefore, planting on fields which are not plowed at all, that is plantings in stubble would fail even worse.

"3. Plowing to a depth of not less than 20 centimeters (about 8 inches) safeguards the field from infestation by weeds. Shallow plowing on the other hand causes serious infestation. Therefore, crops planted in unplowed stubble would be infested even more.

"4. During dry summers, winter crops often suffer from lack of moisture even when planted on good black early fallow. It therefore follows that during dry years crops planted in stubble will die out altogether. This is because the moisture on such fields was not allowed to accumulate but on the contrary was being exhausted by the preceding crops.

"5. Plants feed on minerals. For proper activity of the micro-organisms which mineralize organic substances it is necessary to have air in the soil. In the absence of air there will be no plant food and the plants will starve. Working over the soil produces favorable conditions for the access of air. From this it can be concluded that crops in stubble will certainly starve. On such fields the plants will be small and will produce inferior grain.

"All the explanations brought forth above are, as far as I know, widely accepted among workers in agricultural science.

"Until the fall of 1941, I did not doubt a single one of the accepted explanations of low yields obtained from crops planted in unplowed stubble. However, in the winter of 1941 and spring of 1942 my observations of winter conditions in open unprotected fields led me to a firm conviction that in the Siberian prairies planting on fallow does not present the best conditions for wintering of crops.

"Winter wheat on fallow seldom makes a crop in the Siberian prairies because it is as a rule winter killed. It is for this reason that winter wheat is not being sown in Siberia. Rye if sown at the proper time on good fallow fields will stand the winter but even rye winters poorly or does not come through at all when sown in fields plowed too late. A number of detailed observations led me to suppose that in open sections of Siberia the best winter conditions are assured when winter crops are planted not in fallow but in stubble. At the same time the study of

winter conditions in Siberia led to the conclusion that the favorable winter conditions which exist on fields in stubble can not be produced on large fallow fields.

"By the spring and summer of 1943 it became obvious that the results of the experimental seedings of winter wheat in stubble carried out under my direction on the fields of the Siberian Experimental Institute for Cereal Crops, as well as the data on yields obtained from trial seedings on several state farms, exceeded our expectations. Seedings of the more frost-resistant kinds of winter wheat on good fallow field without snow arresters were entirely lost, which is as should be expected. The highly frost-resistant rye (Omka) which was planted in September was either badly affected or entirely killed. At the same time plantings in stubble of all types of winter wheat, the majority of which were of the Odessa and Crimean types which are of low frost-resisting quality, all came through the winter in excellent condition. Furthermore, on the fields of the Tcheliabinsk station and at Omsk even sprouts of spring wheat wintered over satisfactorily.

"Observations of the growth of plants and data on yields in stubble planting have shown something that we did not suspect when the experiments were started. The results demonstrated that the old, commonly accepted explanation of low yields of rye planted in unplowed stubble in Siberia were not at all valid. The causes of low yields of rye planted in unplowed stubble are entirely different and can be eliminated. Planting in stubble has always been done by the broadcast method. In spite of multiple harrowing, the rye seed was worked into the soil very poorly. The seeds of weeds which require only shallow depth were worked in much better with the harrow than were the seeds of rye. As a result, such broadcast seedings came up too late in the fall. This was especially true under conditions of drought. In the spring and summer of the following year the rye was crowded out by weeds. The horse-drawn drills were also inadequate for working the rye seed into the soil. It is only with tractor-drawn drills that rye can be well worked into the soil. With this method the seeds of weeds which lie on the surface of the ground remain in an unfavorable condition, that is, they are not worked into the soil. Such in-stubble seedings in the prairies of the Siberia not only of winter wheat but also of rye can, as the experiment showed, give yields not lower and in general even higher than the yields of crops sown on fallow and certainly much higher than the entirely unsuitable seedings of rye on fields freshly plowed in September, a practice which unfortunately has recently been widely followed in Siberia. The availability of good tractor-drawn disk drills has made it possible to advocate the old and discarded method of planting in stubble as the better method both agronomically and economically for the prairie regions of Siberia.

"Our proposition is that in the prairie regions of Siberia the planting of winter crops should be carried on in the stubble of spring crops which, in turn, were planted on fallow or sod. Winter wheat must be planted in the last third of August and not later than the first of September. Rye must not be planted later than between the fifth and tenth of September. The crop must be drilled with an adequate tractor-drawn disk drill by the criss-cross method. Tractor-drawn drills with suitable disks and springs can work in the seed properly, while leaving the stubble unbroken and without working in the seed of weeds which lie on the surface. Under Siberian conditions with the great range in day and night temperatures at the end of

August and the middle of September the heavy dew makes possible a stand even with the small amount of precipitation. It is only necessary that the seeding in stubble be done at the proper time with adequate tractor-drawn drills. Crops put in late, that is, in September and especially late in September, can not give a stand even if the crop is put in with tractor-drawn drills and even with good moisture condition. This is because of the low temperature. The results of the tests have already shown that planting in stubble gives the best guarantee against winter killing. It is therefore incorrect to say that stubble plantings of rye in the Siberian prairies are spotty and weak because the plants are killed by frost. The real reason for it is the broadcast method of sowing which, especially when done late, results in a spotty stand and causes the fields to become infested with weeds.

"Both science and experience have unquestionably proved that shallow plowing causes weed infestation. It would naturally appear at first glance that planting of winter crops in unplowed stubble carried on by the method which I propose would cause greater infestation by weeds."


"In my opinion, good stubble seedings of winter wheat and especially of rye made with tractor-drawn disk drills

by the criss-cross method are not only not causing infestation but to some extent appear to be clearing fields from a number of weeds. This does not mean, however, that planting in stubble should be carried out on infested fields. On infested fields no method will give good results.

"It is our deep conviction that the proposed method will in the near future find wide application in the prairie regions of Siberia as a means of obtaining a cheaper supply of winter rye and at the same time improve fertility of the soil for crops that follow. This method of planting will be a useful means for combating soil erosion which, for a number of reasons, is an item of outstanding importance. Finally, the method which we propose makes it possible not only to prevent the loss by winter killing of a large proportion of frost resistant rye, but also opens up wide possibilities for winter wheat."

In the last paragraph Lysenko warns against misapplication of his method. He states:

"I want to emphasize that our proposition applies only to prairie and open timber-prairie regions of Siberia Transural and Northern Kazakhstan. We do not recommend it in other regions, on fields surrounded by forests and therefore protected from wind."



REVIEWS

THE FARMER AND THE REST OF US. By Arthur Moore. Little, Brown & Company, Boston 6, Mass.

This book, by the editor of the Daily Pantagraph of McLean County, Ill., in the heart of the Cornbelt, may not have been intended for the farmers, but certainly many of them will read it with interest. This volume of slightly more than a couple hundred pages is intended for "the rest of us." It is a plea to those of us on paved streets to understand and to begin acting on our most pressing problem, namely, food.

Arthur Moore, the author, is rural enough to recognize the fact that food is fabricated soil fertility, and therefore a product of the farmer's industry. He is also industrial enough to appreciate how we succeeded in bringing on our high production per individual. His broad concepts enable him to discuss the two constituents of American democracy, namely, the farmer and industry, for the elimination of class strife between them and for the good of the cause of food production on the farms. His discussions of their interrelations ought to serve as an awakening experience to any city reader.

The American Farmer's passion for high prices, according to Mr. Moore, has moved farmers to town, has encouraged farm tenancy and has given the Nation its loss of soil fertility, the farmer his loss of security, and all of us the loss of a social democracy. Industrialism has gone forward, but at the cost of the soil body by erosion, as well as of its food-producing power by exhaustion, and of the human resources of the country areas. The cry for cheap food for the rest of us has been an industrial boomerang. It reduced the farmer's capacity to buy industry's products and led us into a depression with its destruction of food and fiber while some people were clamor-

ing for them under Federal aid. The facts, this book points out, demonstrate our need for more than an agricultural economic policy, in the form of a distinctly national food policy.

This book is just what Mr. Moore intended it to be. It is a search "through industrial society for the core of facts about agriculture's relation to the rest of society." It has identified "the thorns and thistles," including both "those which flourish in Washington" and those "which have taken root on the land itself, in the mind of labor, and in industrial management." It analyzes clearly the fallacy in the common belief that "farming is a business" and suggests that "if business would look to agriculture for guidance it might settle its basic controversies, once and for all, in favor of the production man and against the manipulator of finance." In a hungry world profit is not the goal of farming. In terms of food, the farmer can not believe that he produces one hog too many. In terms of economics he is told that he does. Fundamentally, his calling is endowed with more than bookkeeping. It is, Mr. Moore says, "without a counterpart in business, trade or the professions. If we finally hit on a philosophy of farming which will protect our food resources," he says, "it will have to have much of farming's own unique character."

This book puts soil conservation on a new and higher plane, supports this activity in its bidding to become a part of a national food policy. Conservationists of any category will subscribe to his belief that "in soil conservation the people are first and the soil is second," that "we need technicians, but first we need farmers who want technicians," and that "to preserve the land, the farmers—not town-dwellers with their romantic and sentimental illusions—but farmers themselves must believe life on the land to be worth preserving."

The book is another of the recent testimonies that America is now thinking about her soil and the future food production from it. Mr. Moore has thought clearly, written forcefully but sympathetically. His fine style in writing about so universal a concern as food and food production should get a hearing from "the farmer and the rest of us."

—W. A. Albrecht

JUN 29 1945

JUN 23 1945

REFERENCE LIST



Compiled by William L. Robey, Printing and Distribution Unit

SCS personnel should submit requests on Form SCS-37 in accordance with the instructions on the reverse side of the form. Others should address the office of issue.

SOIL CONSERVATION SERVICE

Conservation of Wildlife: Statement of Dr. Edward H. Graham, Chief, Biology Division, Soil Conservation Service, Department of Agriculture; Accompanied by Philip F. Allan, Assistant Chief, Division of Biology, Soil Conservation Service, Department of Agriculture. Division of Biology. Reprint from House Document No. 20. November 27, 1944.

Certain Hydrologic and Climatic Characteristics of the Southwest Region. Regional Bulletin No. 98, Engineering Series No. 9. Soil Conservation Service, United States Department of Agriculture. Region 6. Albuquerque, New Mexico. April 15, 1945. Processed.

Effects of Soil Erosion on Navigation in Upper Chesapeake Bay. Reprinted from The Geographical Review, Vol. XXXV, No. 2. Soil Conservation Service. April, 1945.

Federal-State Cooperative Snow Surveys and Irrigation Water Forecasts for Oregon, April 1, 1945. Division of Irrigation, Soil Conservation Service, United States Department of Agriculture with the cooperation of Oregon Agricultural Experiment Station. April, 1945.

Federal-State Cooperative Snow Surveys and Irrigation Water Forecasts for Utah, April 1, 1945. Division of Irrigation, Soil Conservation Service, United States Department of Agriculture, Utah Agricultural Experiment Station in cooperation with U. S. Forest Service, U. S. Geological Survey, and U. S. National Park Service. April, 1945.

Sediment Complicates Flood Control. Soil Conservation Service, United States Department of Agriculture. Reprinted from Civil Engineering, Vol. 15, No. 2. February, 1945. Processed.

OFFICE OF INFORMATION

U. S. DEPARTMENT OF AGRICULTURE

A Guide to Conservation Practices for Ohio. AIS-18. War Food Administration, Agricultural Adjustment Agency. April, 1945.

Physical Land Conditions in Kent Soil Conservation District, Maryland. Physical Land Survey No. 37. Soil Conservation Service, United States Department of Agriculture. 1945. 15¢. ¹

A Post-War Foreign Trade Program for United States Agriculture. AIS-15. Interbureau Committee on Post-War Programs, United States Department of Agriculture. April, 1945.

Potato Production in the Northeastern and North Central States. Farmers' Bulletin No. 1958. Division of Fruit and Vegetable Crops and Diseases, Bureau of Plant Industry, Soils, and Agricultural Engineering, Agricultural Research Administration. September, 1944. 10¢. ¹

Rural Electrification After the War. AIS-11. Interbureau Committee on Post-War Programs, United States Department of Agriculture. February, 1945.

Some Plain Facts about the Forests. Miscellaneous Publication No. 543. Forest Service, United States Department of Agriculture. April, 1944. Slightly Revised March, 1945. 10¢. ¹

Some Questions . . . and Answers on Where and How to Get a Farm. AIS-19. Bureau of Agricultural Economics, United States Department of Agriculture. April, 1945.

The Work of the U. S. Forest Service. Miscellaneous Publication No. 290. Forest Service, United States Department of Agriculture. January, 1938. Revised January, 1945. 10¢. ¹

STATE BULLETINS

Economic Land Classification in King and Snohomish Counties, Washington, and Its Influence on Full-Time Farm Returns. (Revised). Washington Agricultural Experiment Station, Pullman, Washington. 1944. Processed.

Fifty-Sixth Annual Report. Georgia Experiment Station, Experiment, Georgia. 1943-44. 1945.

Gains on Steers on Native and Reseeded Pasturage: Second Experiment October 13, 1943, to October 31, 1944. Texas Agricultural Experiment Station, A. & M. College of Texas, College Station, Texas with the cooperation of the Soil Conservation Service. Progress Report No. 924. 1945. Processed.

Grain Sorghums Under Irrigation in the Wichita Valley. Progress Report No. 936. Texas Agricultural Experiment Station, A. & M. College of Texas, College Station, Texas. March, 1945.

Important Perennial Weeds in Montana: Their Identification and Control. Bulletin No. 426. Montana State College, Agricultural Experiment Station, Bozeman, Montana. January, 1945.

Irrigated Pastures for Central Washington. Bulletin No. 319. The State College of Washington, Pullman, Washington. January, 1945.

Kaimi Spanish Clover for Humid Lowland Pastures of Hawaii. Circular No. 22. Hawaii Agricultural Experiment Station, Honolulu, U. S. A. February, 1945.

Kudzu in Texas. Progress Report No. 927. Texas Agricultural Experiment Station, A. & M. College of Texas, College Station, Texas. January, 1945.

Michigan's 1945 Crops Program. Bulletin No. F-69. (Revised). Extension Service, Michigan State College, East Lansing, Michigan. January, 1945.

Mississippi Farm Research. Vol. 8, Nos. 1, 2. Mississippi Agricultural Experiment Station, College Station, Mississippi. January, February, 1945.

Montana Insect Pests, 1943 and 1944; Thirtieth Report of the State Entomologist. Bulletin No. 425. Montana State College, Agricultural Experiment Station, Bozeman, Montana. January, 1945.

Plant Propagation. Press Bulletin No. 544. Georgia Agricultural Experiment Station, Experiment Georgia. March, 1945.

¹ From The Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C.

1945



nit

Publi-
part-
vised

ow to
Eco-
ture.

Pub-
part-
uary,

omish
Time
tural
Pro-

ation,

Sec-
1944.
illege
ation
t No.

alley.
peri-
tion,

ifica-
Col-
Mon-

No.
Yash-

es of
peri-

Agri-
exas,

(Re-
East


ssippi
Mis-

rt of
State
man,

Agri-
rgia.

Print-

-643791



S.8 S58a

REFERENCE

SOIL CONSERVATION

Index

VOLUME X

July 1944 to June 1945

**United States Department of Agriculture
Soil Conservation Service
April 1947**



UNITED STATES DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION -- INDEX TO VOLUME X

JULY 1944 TO JUNE 1945

	Page		Page
Alabama, Autauga County, conservation practices by Wadsworth Brothers-----	56-58	BENNETT, H. H.:--Continued	
ALBRECHT, W. A., review of the book, The Farmer and the Rest of Us-----	279	Honored by Costa Ricans-----	246
Alfalfa, successor to poverty grass. C. M. Harsh-----	67-70	Plea for new drainage projects-----	120
American Tree Farms, on West Coast and South, certification and standards-----	250-253	Southern soil conservation platform	231
Arizona, wonder grass. Robert V. Boyle-----	244-246,249	Statement on farmer cooperation with conservation districts-----	135
ARNST, ALBERT:		Bindweed--	
Fighting Fern Without Fire-----	268-270	control by crested wheatgrass.	
Lumber Industry Welcomes Conservation Forestry Farmers-----	250-253	Wilkie Collins, Jr.-----	182-183
Asparagus tops, use as stubble mulch. Otto F. Schnellhardt-----	70-71,72	problem solution by irrigated pastures. Paul G. Moore-----	180-182
Awards--		Biology, land use frontiers. Edward H. Graham-----	193-197
Garden Club of America to H. H. Bennett-----	33	BJARNASON, HAKON: Iceland's Brave Soils Face Tough Opposition-----	84-86
John Deere Medal to C. E. Ramser---	27	Blackland, in Texas, diversification practices. Gordon Webb-----	91-93,110
National Safety Council, to Howard H. Wilson-----	153	Blasting for food. Carl P. Stevens--	9-11
soil conservation, 4-H Club-----	199-201	BLISH, MELVIN: Farm Pond Fights Fire-----	110
BAILEY, R. Y., honor by Kudzu Club of Georgia-----	158	Bonds, war, purchases by Soil Conservation Service employees-----	55
Baltimore Harbor, sedimentation. L. C. Gottschalk-----	11-12	BONNEY, MAURICE C.: Selective Cutting on Cowlitz Project. With Frank B. Harper-----	102-107
BAYS, EARL R.: The Tall Grass is Coming Back-----	163-164	BOYLE, ROBERT V.: Arizona's Wonder Grass-----	244-246,249
Beavers, stocking in Wyoming-----	62	BRINK, WELLINGTON:	
BELL, R. E.: A Business Man Looks At Part-Time Farming. With J. A. Johnston-----	240-243	1944 Ohio Conference-----	50-55
BENNETT, H. H.:		An Earth at Peace with Itself-----	51
Award of Garden Club of America Medal-----	33	Conservation in Cleveland-----	26-27
Departure for South Africa-----	33	Production Increases are Startling-----	99-101
		Britain's colonies, fight against soil erosion. G. M. Roddan-----	216-220
		BROWN, CARL B.: Downstream Interests-----	123-127
		BROWN, GROVER F., review of book: Conservation in the United States----	15
		Farming for Security-----	159

	Page		Page
Buffalo grass--		Conservation--Continued.	
improved outlook for. James		use in growing peanuts. W. M.	
E. Smith, Jr.-----	247-249	Nixon-----	271-272
value for forage and erosion		water-----	207
resistance-----	247-249	See also Soil conservation.	
Business man, opinion on part-time		Cover crop, new, for vineyards.	
farming. R. E. Bell-----	240-243	Homer E. Stennett-----	255
Bussau, Sir A. Louis, visitor from		Cowlitz Farm Forestry Project,	
Australia-----	93	selective cutting. Frank B. Harper	
CAMPBELL, J. PHIL, obituary-----	192	and Maurice C. Bonney-----	102-107
Cannery, new, construction through		Crop production startling increase	
farm forestry. A. C. McIntyre--	107-109	by conservation. Wellington	
Chesapeake Bay area, sedimentation		Brink-----	99-101
problems-----	3-5, 11-12		
Christmas trees, source of added		Deere, John, medal, award to C. E.	
income. Elmer C. Loy-----	197-198	Rammer-----	27
"Clinching up" for the next rough		Dickson, W. F.: Land Utilization	
ride. H. J. Helm-----	187-190, 206	Past...Present...Future. With	
Cleveland, conservation project.		E. G. Grest-----	147-153
Wellington Brink-----	26-27	Diversification, practice on Texas	
Cloudbursts, damage, prevention.		blackland. Gordon Webb-----	91-93, 110
Frank H. Mendell-----	121-122, 127-135	DORIUS, FLOYD W.: More Idaho Spuds	
COHEE, MELVILLE H.: Soil Conserva-		with Green Manure Crops-----	37-38
tion Districts and Returning		Downstream interests. Carl B.	
Veterans-----	235-238	Brown-----	123-127
COLLINS, WILKIE JR.:		Drainage--	
Crested Wheatgrass Helps Control		problems in Wisconsin districts----	229
Bindweed-----	182-183	projects, prospects for ex-service-	
Irrigated Mountain Meadows-----	165-167	men-----	120
COMES, L. R.: Many Groups Merge		See also Irrigation.	
Efforts in Leelanau-----	34-36, 39		
Contractors, discovery of soil		Earth at peace with itself. Wellin-	
conservation districts. A. E.		ton Brink-----	51
McClymonds-----	109-110	ECKLES, W. F., redesign of farm.	
Conservation--		Barrington King-----	94-98
farmers in forestry, cooperation		ELLISON, W. D.:	
with lumber industry. Albert		Raindrops, Surface Flow and	
Arnst-----	250-253	Erosion-----	128-129
in Cleveland. Wellington Brink--	26-27	When Raindrops Splash-----	58-62
practices in Alabama by Wadsworth		ENLOW, C. R.: Legume and Grass	
Brothers. Barrington King-----	56-58	Seed Production 1944-----	118-120
progress report in Wyoming. A. E.		Erosion--	
McClymonds-----	203	effect of raindrops. W. D.	
rides timber trail in Turtle Moun-		Ellison-----	128-129
tains. Walter H. Paul and Elmer		effects on plant and animal com-	
L. Worthington-----	13-15	munities-----	197
soil. See Soil conservation		soil--	
total, greater task ahead. R. W.		control--	
Rogers-----	202-203	postwar program in Wisconsin 228-230	
		problem in Britain's colonies 216-220	

	Page
Erosion--Continued.	
soil--Continued.	
effects of downstream enter-	
prises-----	123-127
from cloudbursts, prevention-	121-122,
	127,135
problems in Iceland-----	81-83,86-87
splash, causes and occurrence on	
dry soil-----	58-62
Ethiopia, agricultural survey,	
A. T. Semple-----	154-157
Europe's needs, report-----	270
Explosives, use in blasting swamp	
areas for pasture sites-----	9-11
Farm--	
forestry. See Forestry, farm	
importance of conservation research.	
R. E. Uhland-----	43-49
pond. See Pond, farm	
redesign by W. F. Eckles. Barrington	
King-----	94-98
Farmers, cooperation with soil con-	
servation districts-----	135
Farming, part-time, outlook for a	
business man. R. E. Bell-----	240-243
Farms--	
damage when Maumee River runs brown.	
Thomas H. Langlois-----	6-8,15
fertilizing for fertility,	
maintenance-----	253-254
FERGUSON, E. T.: Gully Head Control	
in Deep Loess-----	129
Fern, fighting without fire. Albert	
Arnst-----	268-270
Fertility level of soils, raising.	
B. H. Hendrickson-----	253-254
Field border ecology-----	195
FIERO, KENNETH: Districts Ride the	
Range-----	144-146
Fighting fern without fire. Albert	
Arnst-----	268-270
Fire--	
control by water from farm ponds.	
Melvin Blish-----	110
disadvantage in fighting fern.	
Albert Arnst-----	268-270
Fires, in woods outlaw by county.	
L. J. Leffelman-----	157

	Page
Fish, damage by slit in Maumee River.	
Thomas M. Langlois-----	6-8,15
FISHEL, RAYMOND N.: Two Home-Made	
Tree Planters. With Marion M.	
Weaver-----	71-72
Flood control problems in Wisconsin	
districts-----	229
Food, blasting for. Carl P. Stevens-	9-11
Forage, abundance in Northern Great	
Plains-----	187-190,206
Forestry--	
conservation, farmers, cooperation	
with lumber industry. Albert	
Arnst-----	250-253
farm--	
benefits from selective cut-	
ting-----	102-107
projects in time of war. John	
W. Keller-----	221-224
source of lumber for new cannery.	
A. C. McIntyre-----	107-109
to the rescue. Frank B. Harper	
and Irving R. Pearce-----	77-78,88
FORT T. HICKS, JUDGE, statement to	
Taylor County, Ga. Grand Jury-----	29,33
4-H Club, national awards in soil	
conservation. Walter W. John---	199-201
Frontiers in land use biology.	
Edward H. Graham-----	193-197
Garden Club of America, award to	
H. H. Bennett-----	33
GETZE E. BLOREN, JR.: Bill Roper,	
Lancaster Leader-----	215,227
GLOVER, FRANKLIN M., JR.: Harvesting	
Seed of Barbados Sourgrass-----	62-63
GOITSCHALK, L. C.: Sedimentation in	
a Great Harbor-----	3-5,11-12
GRAHAM, EDWARD H.:	
Frontiers in Land Use Biology---	193-197
Review of book Practical Farming	
for the South-----	63-64
Grass--	
poverty, replacement by alfalfa.	
C. M. Harsh-----	67-70
seed--	
production, 1944. C. R. Enlow	118-120
shortage, solution-----	8
tall, return in Colorado. Earl R. Bays	163-164

	Page
Grassland, improvement for better livestock. R. H. Lush-----	272-274
Grazing facilities--	
improvement in Northern Great Plains-----	187-190,205
See also Range.	
Great Plains, shelterbelts. John W. Keller-----	259-263
Green manure, use for increasing Idaho potato yields. Floyd W. Dorius-----	37-38
QUEST, E. G.: Land Utilization Past...Present...Future. With W. F. Dickerson-----	147-153
Gullies control in deep loess by flumes. E. T. Ferguson-----	129
Gully problems in Wisconsin districts	229
HAMILTON, J. G.: Irrigated Pastures are a "Money Crop."-----	176-178
HARPER, FRANK B.: Farm Forestry to the Rescue. With Irving F. Pearce-----	77-78,88
Iceland's Son Works on Problem of Shifting Sands-----	81-83,86-87
Irrigated Pastures Spurred by War. With Harold E. Tower-----	172-175
Selective Cutting on Cowlitz Project. With Maurice C. Bonney--	102-107
HARSH, C. M.: From Poverty Grass to Alfalfa-----	67-70
Harvesting seed of Barbados Sourgrass. Franklin M. Glover, Jr.--	62-63
Hedge management, research-----	195
HELM, J. J.: "Cinching Up" for the Next Rough Ride-----	187-190,205
HENDRICKSON, B. H.: Raising the Fertility Level-----	253-254
HOLLINGER, EDWIN C., review of book, Biology for Better Living-----	111
HUGHES, R. R., designation as "Master Conservationist"-----	146
Iceland--	
agriculture, description----	81-83,86-87
farm boy, study of problem of shifting sands. Frank B. Harper----	81-83,86-87
soils, fight against erosion. Hakon Bjarnason-----	84-86

	Page
Income, additional, from Christmas trees. Elmer C. Loy-----	197-198
Indiana, Switzerland County, referendum methods-----	25
Insect damage and control-----	196
Irrigation--	
of pastures--	
increase by War. Harold E. Tower and Frank B. Harper-----	172-175
solution to bindweed problem. Paul G. Moore-----	180-192
source of "money crop". J. G. Hamilton-----	176-178
on mountain meadows. Wilkie Collins, Jr.-----	165-167,175
value of water-spreading formulae-----	73-76,87
See also Drainage.	
Job No. 1, postwar: Soil Conservation. Claude R. Wickard-----	19-23
JOHN, WALTER W.: Meet Jim Skinner of Michigan----	239,254
National 4-H Awards in Soil Conservation-----	199-201
JOHNSON, J. A.: A Business Man Looks at Part-Time Farming, with R. E. Bell-----	240-243
KELLER, JOHN W.: Farm Forestry Projects in Time of War-----	221-224
Shelterbelts in the Great Plains	259-263
KING, BARRINGTON: Eckles Redesigns His Farm-----	94-98
The Wadsworth Brothers, a Conservation Team-----	56-58
KRIMGOLD, B. B.: Translation of article by Academician T. Lysenko on Planting of Winter Wheat and Rye in Unplowed Stubble-----	277-279
Kudzu--	
crowns, disposal through advertising-trial at Beltsville, Md.-----	164
Kudzu Club of Georgia-----	158
KUNZE, GLADYS: Soil Conservation--Its Relationship to Wildlife and Human Welfare-----	28-29,33

	Page
Land--	
exchange for water through conser- vation districts. John G.	
Sutton-----	130-135
use biology frontier. Edward H.	
Graham-----	193-197
utilization--	
past, present, and future, E. G.	
Grest and W. F. Dickson-----	147-153
projects, restoration of Old West. O. E. McConnell-----	264-267
Lands, submarginal, in Wisconsin districts-----	229
LANGLOIS, THOMAS H.: Fish and Farms Suffer When the Maumee Runs Brown 6-8,15	
LANTOW, J. L.: Plant Vigor and Range Production-----	178-180
LATHROP, WILLIAM H.: Seeing is Believing-----	24-25
Leelanau County, Michigan, group efforts for conservation. L. R. Combs-----	34-36,39
LEFFELMAN, L. J.: County Outlaws Woods Fires-----	157
Legume seed--	
production, 1944. C. R. Enlow--	118-120
shortage, solution-----	8
LEOPOLD, ALDO, review of book, Natural Principles of Land Use-----	38-39
Lespedeza seed, planting in South Carolina districts-----	229
Lime production problems in Wisconsin districts-----	299
Livestock, improvement through better grassland. R. H. Lush-----	272-274
Loess, deep, need for gully head control. E. T. Ferguson-----	192
Logging by selective cutting, Cowlitz Farm Forestry Project. Frank E. Harper and Maurice C. Bonney----	102-107
Lovegrass Lehman, wonder of Arizona. Robert V. Boyle-----	244-246,249
LOY, ELMER C.: Added Income from Christmas Trees-----	197-198
LOYD, GLENNON: Terracing for Fun	115-117
Lumber industry, cooperation with conservation forestry farmers. Albert Arnst-----	250-253
LUSH, R. H.: Improve Grassland for Better Livestock-----	272-274

	Page
LYSENKO, T.: Article, on planting of Winter Wheat and Rye in Unplowed Stubble. Translated and abstracted by D. B. Kringold-----	277-279
Machinery, potato, improvement by recent research. John W. Slosser-----	191-192
Marsh management for muskrat pro- duction-----	194
"Master Conservationist," designation of. R. R. Hughes-----	146
MCCALLA, T. M.: Microbiology and Soil Conservation-----	225-227
MCCLYMONDS, A. E.: Account of Stewardship-----	203
Contractors Discover Soil Conserva- tion Districts-----	109-110
MCCONNELL, O. E.: Old West Lives Again in LU Projects-----	264-267
MCINTYRE, A. C.: Farm Forestry Builds a New Cannery-----	107-109
Meadows, mountain, irrigation. Wilkie Collins, Jr.-----	165-167,175
Memorials, woodlands for-----	158-159
MEDELL, FRANK H.: Gone with the Cloudburst-----	121-122,127,135
Michigan--	
farmer, Jim Skinner. Walter W. John-----	239,254
Leelanau County, group organiza- tions for conservation. L. R. Combs-----	34-36,39
muckland, reed canary grass pro- fit. W. L. Williams-----	204-205
Microbiology and soil conservation. T. M. McCalla-----	225-227
Midwest Plow Terrace Building Con- test-----	115-117
MILES, WAYNE H.: Water Spreading	73-76,87
MOHAGEN, VERNA C.: Review of book-- Bureaucracy, A Challenge To Better Management-----	110-111
Trampling out the Vintage-----	230-231
They Scout the World-----	139-143,146
MOORE, PAUL G.: Irrigated Pastures Solve Bindweed Problem-----	180-182
Mountain meadows, irrigation. Wilkie Collins, Jr.-----	165-167,175

	Page
Muckland, Michigan, reed canary grass profits. W. L. Williams-----	204-205
Mulch, stubble, use of asparagus tops. Otto F. Schnellhardt-----	70-71, 72
Muskrats, dividends from, on farm near Worland, Wyo.-----	62
National 4-H Club Soil Conservation Contest, scholarship winners-----	199-201
National Safety Council award to Howard W. Wilson-----	153
NEAL, O. R.: Water Storage in the Surface Soil-----	274-277
Neglect, example in South Carolina. Jim Fort Rast-----	79-80
1944 Ohio Conference. Wellington Brink-----	50-55
NIXON, W. M.: Growing Peanuts the Conservation Way-----	271-272
Northern Great Plains, range management improvement-----	187-190, 206
Ohio--	
conference on conservation, nutrition, and human health. Wellington Brink-----	50-55
Coshocton County, soil conservation program-----	67-70
Oklahoma, benefits from good pastures-----	211-214
Old West, restoration by LU projects. O. E. McConnell-----	264-267
Pastures--	
good source of increased gains and profits. H. W. Staten-----	211-214
irrigation--	
increase by War. Harold E. Tower and Frank B. Harper-----	172-175
solution to bindweed problem. Paul G. Moore-----	180-182
value as "money crop." J. G. Hamilton-----	176-178
PAUL, WALTER H.: Conservation Rides Timber Trail in Turtle Mountains. With Elmer L. Worthington-----	13-15
Peanuts, growing the conservation way. W. M. Nixon-----	271-272
PEARCE, IRVING F.: Farm Forestry to the Rescue. With Frank B. Harper-----	77-78, 88

	Page
Pennsylvania, Lancaster County, soil conservation work of Hartwell E. Roper-----	215-227
Plant vigor and range production. J. L. Lantow-----	178-180
Planters, tree, two-home-made. Marion M. Weaver and Raymond N. Fishel-----	71-72
Plants, noxious, problems and control-----	196-197
Pond, farm--	
management for fish production- value in fire fighting. Melvin Blish-----	194-195
Postwar job No. 1: Soil Conservation. Claude R. Wickard-----	110
Potato machinery, improvement by recent research. John W. Slosser- Potatoes, Idaho, increases with green manure-----	191-192
PRESTON, JOHN F., review of books: Woodland Opportunities on Dairy Farms in New York-----	37-38
Some Ecological Aspects of Afforestation and Forestry in Great Britain	87
Profits, result of range care. W. T. White-----	206
Progressive Farmer, tribute to H. H. Bennett-----	168-171, 175
231	
Raindrops--	
cause of surface flow and erosion. W. D. Ellison-----	128-129
splashing, consequences. W. D. Ellison-----	58-62
RAMSER, C. E., award of John Deere Medal-----	27
Range--	
assistance by soil conservation districts. Kenneth Fiero-----	144-146
land and adjustments, making in Utah C. P. Starr and D. C. Winn-----	30-31
lands, conservation practices and profits-----	168-171, 175
management, improvement in Northern Great Plains-----	187-190, 206
production and plant vigor. J. L. Lantow-----	178-180
profits through range care. W. T. White-----	168-171, 175

	Page
RAST, JIM FORT: There Is No Sadder Sight-----	79-80
Reed canary grass profits on Michigan muckland. W. L. Williams-----	204-205
Reforestation in Wisconsin districts- Research--	229
grows up on the farm. R. E.	
Uhland-----	43-49
recent on potato machinery improve- ment. John W. Slosser-----	191-192
Reviews--	
Bureaucracy: A Challenge to Better Management. J. M. Juran-----	110-111
Biology for Better Living. Ernest C. Bayles and F. Will Burnett---	111
Conservation in the United States. A. F. Gustafson, C. H. Guise, W. J. Hamilton, Jr., and H. Ries-----	15
Farming for Security. William E. Duryee-----	159
Natural Principles of Land Use. Edward H. Graham-----	38-39
Practical Farming for the South. Benjamin F. Bullock-----	63-64
Some Ecological Aspects of Afforestation and Forestry in Great Britain. Sir Roy Robinson-----	206
The Farmer and the Rest of Us. Arthur Moore-----	279
The Nature and Properties of Soils. T. Lyttleton Lyon and Harry O. Buckman-----	205
Trampling Out the Vintage. Joseph A. Cocannouer-----	230-231
Woodland Opportunities on Dairy Farms in New York. Hugh A. Johnson, Irving C. Fellows, Donald Rush, C. R. Lockard, and C. Edward Behre-	87
RIST, BOYD, banker, statement on soil conservation-----	32
ROBEV, WILLIAM I.: Soil conservation bibliographies. See each issue July-June, incl.	
RODDAN, G. M.: Fighting Soil Erosion in Britain's Colonies-----	216-220
Rodents, wild, biological control----	196
ROGERS, R. W.: The Greater Task Ahead-----	202-203
Roper, Bill, Lancaster leader. E. Bioren Getze, Jr.-----	215-227

	Page
Rye, planting in unplowed stubble, article by Academician T. Lysenko. Translated and abstracted by D. B. Krimgold-----	277-293
Sands, shifting, problems, study by Paul Sveinsson of Iceland. Frank B. Harper-----	81-83, 86-87
SCHNELLHARDT, OTTO F.: Stubble Mulching with Asparagus Tops--	70-71, 72
Sedimentation in Baltimore harbor. L. C. Gottschalk-----	3-5, 11-12
Seed harvesting of Barbados sourgrass. Franklin M. Glover, Jr.-----	62-63
Seeing is believing. William H. Lathrop-----	24-25
SEMPLE, A. T.: A Look at Ethiopia Shelterbelts--	154-157
Great Plains. John W. Keller---	259-263
increases production in Texas-----	25
SHIFFER, ORRIE: Wisconsin Districts Consider Post-War Period-----	228-230
Skinner, Jim, from Michigan. Walter W. John-----	239-254
SLOSSER, JOHN W.: Recent Research Improves Potato Machinery-----	191-192
SMITH, JAMES E., JR.: Outlook Improves for Buffalo Grass-----	247-249
Soil--	
conservation--	
and microbiology. T. M. McCalla-----	225-227
conference, 1944, in Ohio-----	50-55
districts--	
and returning veterans. Melville H. Cohee-----	235-238
discovery by contractors. A. E. McClymonds-----	109-110
exchange of water for land. John G. Sutton-----	130-135
organization in Michigan, Lee-lanau County-----	34-36, 39
postwar period, outlook in Wisconsin. Orrie Shiffer-	228-230
referendum methods, Indiana, Clark and Switzerland Counties-----	25
ride the range. Kenneth Fiero-----	144-146
work, expansion-----	80

	Page		Page
Soil--Continued.		Stone fence projects in Wisconsin	
conservation--Continued.		districts-----	229
in uplands, effects on downstream		Storage, of water in surface soil.	
enterprises-----	123-127	O. R. Neal-----	274-277
national 4-H awards. Walter W.		Streambank erosion, Wisconsin	
John-----	190-201	districts-----	229
No. 1 postwar job. Claude R.		Streamside ecology and erosion con-	
Wickard-----	19-23	trol-----	194-195
relationship to wildlife and human		Stubble--	
welfare. Gladys Kunze-----	28-29,33	mulching with asparagus tops. Otto F.	
role, in privately-owned range lands		Schnellhardt-----	70-71,72
of Pacific northwest-----	168-171,175	unplowed, planting of winter wheat	
statement by--		and rye in article by Academician T.	
Boyd Rist, banker-----	32	Lysenko. Translated and abstracted	
Judge Fort-----	29,33	by D. B. Kringold-----	277-293
See also Conservation.		Summers, George E., demonstration of	
conservationists scout the world.		contour farming-----	24-25
Verna C. Mohagen-----	129-143,146	Surface--	
erosion. See Erosion, soil.		flow, effect of raindrops. W. D.	
fertility level, raising. B. H.		Ellison-----	128-129
Hendrickson-----	253-254	soil, effects of water storage.	
surface, effects of water storage.		O. R. Neal-----	274-277
O. R. Neal-----	274-277	SUTTON, JOHN G.: Districts Exchange	
Soil Conservation Service, research		Water for Land-----	130-135
program-----	43-49	Swamp areas, blasting for pasture	
Soils--		sites in Washington-----	9-11
microbiology and conservation. T. M.		Terracing for fun. Glennon Loyd-	115-117
McCalla-----	225-227	Texas blackland, profits from diversi-	
of Iceland, fight against erosion.		fication. Gordon Webb-----	91-93,110
Hakon Bjarnason-----	84-86	Timber trail, in Turtle Mountains,	
Sourgrass, Barbados, seed, harvesting.		application of conservation to.	
Franklin M. Glover, Jr.-----	62-63	Walter R. Paul and Elmer L.	
South Africa, Union of, invitation to		Worthington-----	13-15
H. H. Bennett-----	33	TOWER, HAROLD E.: Irrigated Pastures	
Southern soil conservation platform.		Spurred by War. With Frank B.	
H. H. Bennett-----	231	Harper-----	172-175
Splash erosion, causes and occurrence		Tree--	
on dry soil-----	58-62	farm system, description and	
Spoilbanks, management, research--	195-196	operation-----	250-253
Spuds, Idaho, increases with green		planters, two home-made. Marion M.	
manure crops. Floyd W. Dorius-----	37-38	Weaver and Raymond N. Fishel----	71-72
STARR, C. P.: Making Range Adjustments		Trees,-Christmas, source of added	
in Utah. With D. S. Winn-----	30-31,39	income. Elmer C. Loy-----	197-198
STATEN, Hi W.: Increased Gains and		TUCKER, R. J., statement on value of	
Profits Through Good Pasture-----	211-214	shelterbelts in Texas-----	25
STEELE, J. GORDON, review of book,		Turtle Mountains, timber trail in	
The Nature and Properties of Soils	205	application of conservation. Walter	
STENNETT, HOMER E.: A New Cover Crop		H. Paul and Elmer L. Worthington--	13-15
for Vineyards-----	255		
STEVENS, CARL P.: Blasting for Food	9-11		

	Page		Page
UHLAND, R. E.: Research Grows Up on the Farm-----	43-49	WEAVER, MARION M.: Two Home-Made Tree Planters. With Raymond N. Fishel-----	71-72
Union of South Africa, invitation to H. H. Bennett-----	33	WEBB, GORDON: Diversification Comes to Texas Blackland-----	91-93, 110
Utah, range adjustments in. C. P. Starr and D. S. Winn-----	30-31, 39	Welfare, human, relations to soil conservation. Gladys Kunze---	28-29, 33
Vermont Otto Creek Soil Conservation District, report, 1941-43, excerpts-----	99-101	West Coast Tree Farms, certification, standards-----	250-253
Veterans--		Wheat, winter, planting in unplowed stubble, article by Academician T. Lysenko. Translated and abstracted by D. B. Krimgold-----	277-293
employment on drainage projects----	120	Wheatgrass, crested, value in control of bindweed. Wilkie Collins, Jr.-----	182-183
returning, aid by soil conservation districts. Melville H. Cohee- 235-238		WHITE, W. T.: Take Care of the Ranges and the Profits Take Care of Themselves-----	168-171, 175
Vineyards, new cover crop for. Homer E. Stennett-----	255	WICKARD, CLAUDE R.: Soil Conservation: No. 1 Postwar Job-----	19-23
Wadsworth Brothers, a conservation team. Barrington King-----	56-58	Wildlife, relation to soil conservation. Gladys Kunze-----	28-29, 33
Wallace Asa B., Memorial Essay Contest winner-----	28	WILLIAMS, W. L.: Reed Canary Pays Off on Michigan Muckland-----	204
War, effect on irrigation of pastures. Harold E. Tower and Frank B. Harper-----	172-175	Wilson, Howard H., 1944 National Safety Council Award-----	153
War Bonds purchases by Soil Conservation Service employees-----	55	WINN, D. S.: Making Range Adjustments in Utah. With C. P. Starr-----	30-31, 39
Wartime, farm forestry projects. John W. Keller-----	221-224	Wisconsin districts, outlook for post-war period. Orrie Shiffer-----	228-230
Washakie Conservation District, Wyo., progress report-----	203	Woodland--	
Washington, Cowlitz Farm Forestry Project, selective cutting on-----	102-107	products, harvesting for wartime needs-----	77-78, 88
Water--		resources, use in rebuilding cannery-----	107-109
conservation-----	207	Woodlands for memorials-----	158-159
exchange for land through conservation districts. John G. Sutton----	130-135	Woods, fires; outlaw by county. L. J. Leffelman-----	157
spreading. Wayne H. Miles----	73-76, 87	WORTHINGTON, ELMER, L.: Conservation Rides Timber Trail in Turtle Mountains. With Walter H. Paul	13-15
storage in--			
the surface soil. O. R. Neal- 274-277			
Wisconsin districts-----	229		

